

stn

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal612bxr

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page for STN Seminar Schedule - N. America
NEWS 2 JUL 28 CA/CAPLUS patent coverage enhanced
NEWS 3 JUL 28 EPFULL enhanced with additional legal status
information from the EPOline Register
NEWS 4 JUL 28 IFICDB, IFIPAT, and IFIUDB reloaded with enhancements
NEWS 5 JUL 28 STN Viewer performance improved
NEWS 6 AUG 01 INPADOCDB and INPAFAMDB coverage enhanced
NEWS 7 AUG 13 CA/CAPLUS enhanced with printed Chemical Abstracts
page images from 1967-1998
NEWS 8 AUG 15 CAOLD to be discontinued on December 31, 2008
NEWS 9 AUG 15 CAPLUS currency for Korean patents enhanced
NEWS 10 AUG 27 CAS definition of basic patents expanded to ensure
comprehensive access to substance and sequence
information
NEWS 11 SEP 18 Support for STN Express, Versions 6.01 and earlier,
to be discontinued
NEWS 12 SEP 25 CA/CAPLUS current-awareness alert options enhanced
to accommodate supplemental CAS indexing of
exemplified prophetic substances
NEWS 13 SEP 26 WPIDS, WPINDEX, and WPIX coverage of Chinese and
and Korean patents enhanced
NEWS 14 SEP 29 IFICLS enhanced with new super search field
NEWS 15 SEP 29 EMBASE and EMBAL enhanced with new search and
display fields
NEWS 16 SEP 30 CAS patent coverage enhanced to include exemplified
prophetic substances identified in new Japanese-
language patents
NEWS 17 OCT 07 EPFULL enhanced with full implementation of EPC2000
NEWS 18 OCT 07 Multiple databases enhanced for more flexible patent
number searching
NEWS 19 OCT 22 Current-awareness alert (SDI) setup and editing
enhanced
NEWS 20 OCT 22 WPIDS, WPINDEX, and WPIX enhanced with Canadian PCT
Applications
NEWS 21 OCT 24 CHEMLIST enhanced with intermediate list of
pre-registered REACH substances

NEWS EXPRESS JUNE 27 08 CURRENT WINDOWS VERSION IS V8.3,
AND CURRENT DISCOVER FILE IS DATED 23 JUNE 2008.

Updated Search

stn

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items
NEWS IPC8 For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 06:48:32 ON 27 OCT 2008

=> file casreact		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'CASREACT' ENTERED AT 06:48:45 ON 27 OCT 2008
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications.

FILE CONTENT:1840 - 25 Oct 2008 VOL 149 ISS 18

New CAS Information Use Policies, enter HELP USAGETERMS for details.

*
* CASREACT now has more than 15.3 million reactions *
*

CASREACT contains reactions from CAS and from: ZIC/VINITI database (1974-1999) provided by InfoChem; INPI data prior to 1986; Biotransformations database compiled under the direction of Professor Dr. Klaus Kieslich; organic reactions, portions copyright 1996-2006 John Wiley & Sons, Ltd., John Wiley and Sons, Inc., Organic Reactions Inc., and Organic Syntheses Inc. Reproduced under license. All Rights Reserved.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=>
Uploading C:\Documents and Settings\brobinson1\My Documents\e-Red Folder\10524345\365.str

Updated Search

stn

L1 STRUCTURE UPLOADED

=> s l1, crd ibib abs, fh1t,
2540 'L1'
2 CRD
0 IBIB
16097 ABS
3 FHIT

L2 0 L1, CRD IBIB ABS, FHIT,
('L1' (W) CRD (W) IBIB (W) ABS (W) FHIT)

=> d l1, ibib abs crd, fhits

L1 HAS NO ANSWERS

'IBIB ABS CRD FHITS ' IS NOT A VALID STRUCTURE FORMAT KEYWORD
ENTER STRUCTURE FORMAT (SIM), NOS:end

=> d l1, ibib abs crd, fhitr

L1 HAS NO ANSWERS

'IBIB ABS CRD FHITSR ' IS NOT A VALID STRUCTURE FORMAT KEYWORD
ENTER STRUCTURE FORMAT (SIM), NOS:end

=> s l1

SAMPLE SEARCH INITIATED 06:58:26 FILE 'CASREACT'
SCREENING COMPLETE - 179 REACTIONS TO VERIFY FROM 35 DOCUMENTS

100.0% DONE 179 VERIFIED 0 HIT RXNS 0 DOCS
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED VERIFICATIONS: 2778 TO 4382
PROJECTED ANSWERS: 0 TO 0

L3 0 SEA SSS SAM L1 (0 REACTIONS)

=> s l1 full

THE ESTIMATED SEARCH COST FOR FILE 'CASREACT' IS 117.50 U.S. DOLLARS
DO YOU WANT TO CONTINUE WITH THIS REQUEST? (Y)/N or END:y
FULL SEARCH INITIATED 06:58:30 FILE 'CASREACT'
SCREENING COMPLETE - 5687 REACTIONS TO VERIFY FROM 617 DOCUMENTS

100.0% DONE 5687 VERIFIED 70 HIT RXNS 23 DOCS
SEARCH TIME: 00.00.02

L4 23 SEA SSS FUL L1 (70 REACTIONS)

=> d l4, ibib abs fhitr, 1-23

'FHITR' IS NOT A VALID FORMAT FOR FILE 'CASREACT'

The following are valid formats:

ABS ----- GI and AB
ALL ----- BIB, AB, IND, RE, Single-step Reactions
APPS ----- AI, PRAI
BIB ----- AN, plus Bibliographic Data
CAN ----- List of CA abstract numbers without answer numbers

Updated Search

stn

CBIB ----- AN, plus Compressed Bibliographic Data
DALL ----- ALL, delimited (end of each field identified)
IABS ----- ABS, indented with text labels
IALL ----- ALL, indented with text labels
IBIB ----- BIB, indented with text labels
IND ----- Indexing data
IPC ----- International Patent Classifications
ISTD ----- STD, indented with text labels
OBIB ----- AN, plus Bibliographic Data (original)
OIBIB ----- OBIB, indented with text labels

SBIB ----- BIB, no citations
SIBIB ----- IBIB, no citations

MAX ----- Same as ALL
PATS ----- PI, SO
SCAN ----- TI and FCRD (random display, no answer number. SCAN
must be entered on the same line as DISPLAY, e.g.,
D SCAN.)
SSRX ----- Single-Step Reactions (Map, Diagram, and Summary for
all single-step reactions)
STD ----- BIB, IPC, and NCL

CRD ----- Compact Display of All Hit Reactions
CRDREF ----- Compact Reaction Display and SO, PY for Reference
FHIT ----- Reaction Map, Diagram, and Summary for first
hit reaction
FHITCBIB --- FHIT, AN plus CBIB
FCRD ----- First hit in Compact Reaction Display (CRD) format
FCRDREF ----- First hit in Compact Reaction Display (CRD) format with
CA reference information (SO, PY). (Default)
FSPATH ----- PATH, plus Reaction Summary for the "long path"
FSPATH ----- SPATH, plus Reaction Summary for the "short path"
HIT ----- Reaction Map, Reaction Diagram, and Reaction
Summary for all hit reactions and fields containing
hit terms
OCC ----- All hit fields and the number of occurrences of the
hit terms in each field. Includes total number of
HIT, PATH, SPATH reactions. Labels reactions that have
incomplete verifications.
PATH ----- Reaction Map and Reaction Diagram for the "long
path". Displays all hit reactions, except those
whose steps are totally included within another hit
reaction which is displayed
RX ----- Hit Reactions (Map, Diagram, Summary for all hit reactions)
RXG ----- Hit Reaction Graphics (Map and Diagram for all hit reactions)
RXL ----- Hit Reaction Long (Map, Diagram, Summary for all hit reactions)
RXS ----- Hit Reaction Summaries (Map and Summary for all hit reactions)
SPATH ----- Reaction Map and Reaction Diagram for the "short
path". Displays all single step reactions which
contain a hit substance. Also displays those
multistep reactions that have a hit substance in both
the first and last steps of the reaction, except for
those hit reactions whose steps are totally included
within another hit reaction which is displayed

stn

To display a particular field or fields, enter the display field codes. For a list of the display field codes, enter HELP DFIELDS at an arrow prompt (=>). Examples of combinations include: D TI; D BIB RX; D TI, AU, FCRD. The information is displayed in the same order as the specification. All of the formats, except CRD, CRDREF, FHIT, PATH, FPATH, SPATH, FSPATH, FCRD, FCRDREF, HIT, RX, RXG, RXS, SCAN, and OCC, may be used with the DISPLAY command to display the record for a specified Accession Number.

ENTER DISPLAY FORMAT (FCRDREF):end

=> d his

(FILE 'HOME' ENTERED AT 06:48:32 ON 27 OCT 2008)

FILE 'CASREACT' ENTERED AT 06:48:45 ON 27 OCT 2008

```
L1      STRUCTURE UPLOADED
L2      0 S L1, CRD IBIB ABS, FHIT,
L3      0 S L1
L4      23 S L1 FULL
```

=> d l4, ibib abs fhistr, 1-5

'FHISTR' IS NOT A VALID FORMAT FOR FILE 'CASREACT'

The following are valid formats:

```
ABS ----- GI and AB
ALL ----- BIB, AB, IND, RE, Single-step Reactions
APPS ----- AI, PRAI
BIB ----- AN, plus Bibliographic Data
CAN ----- List of CA abstract numbers without answer numbers
CBIB ----- AN, plus Compressed Bibliographic Data
DALL ----- ALL, delimited (end of each field identified)
IABS ----- ABS, indented with text labels
IALL ----- ALL, indented with text labels
IBIB ----- BIB, indented with text labels
IND ----- Indexing data
IPC ----- International Patent Classifications
ISTD ----- STD, indented with text labels
OBIB ----- AN, plus Bibliographic Data (original)
OIBIB ----- OBIB, indented with text labels

SBIB ----- BIB, no citations
SIBIB ----- IBIB, no citations

MAX ----- Same as ALL
PATS ----- PI, SO
SCAN ----- TI and FCRD (random display, no answer number. SCAN
must be entered on the same line as DISPLAY, e.g.,
D SCAN.)
SSRX ----- Single-Step Reactions (Map, Diagram, and Summary for
all single-step reactions)
STD ----- BIB, IPC, and NCL

CRD ----- Compact Display of All Hit Reactions
CRDREF ----- Compact Reaction Display and SO, PY for Reference
```

Updated Search

stn

FHIT ----- Reaction Map, Diagram, and Summary for first
hit reaction
FHITCBIB --- FHIT, AN plus CBIB
FCRD ----- First hit in Compact Reaction Display (CRD) format
FCRDREF ---- First hit in Compact Reaction Display (CRD) format with
CA reference information (SO, PY). (Default)
FPATH ----- PATH, plus Reaction Summary for the "long path"
FSPATH ----- SPATH, plus Reaction Summary for the "short path"
HIT ----- Reaction Map, Reaction Diagram, and Reaction
Summary for all hit reactions and fields containing
hit terms
OCC ----- All hit fields and the number of occurrences of the
hit terms in each field. Includes total number of
HIT, PATH, SPATH reactions. Labels reactions that have
incomplete verifications.
PATH ----- Reaction Map and Reaction Diagram for the "long
path". Displays all hit reactions, except those
whose steps are totally included within another hit
reaction which is displayed
RX ----- Hit Reactions (Map, Diagram, Summary for all hit reactions)
RXG ----- Hit Reaction Graphics (Map and Diagram for all hit reactions)
RXL ----- Hit Reaction Long (Map, Diagram, Summary for all hit reactions)
RXS ----- Hit Reaction Summaries (Map and Summary for all hit reactions)
SPATH ----- Reaction Map and Reaction Diagram for the "short
path". Displays all single step reactions which
contain a hit substance. Also displays those
multistep reactions that have a hit substance in both
the first and last steps of the reaction, except for
those hit reactions whose steps are totally included
within another hit reaction which is displayed

To display a particular field or fields, enter the display field
codes. For a list of the display field codes, enter HELP DFIELDS
at an arrow prompt (=>). Examples of combinations include: D TI;
D BIB RX; D TI, AU, FCRD. The information is displayed in the same order
as the specification. All of the formats, except CRD, CRDREF, FHIT, PATH,
FPATH, SPATH, FSPATH, FCRD, FCRDREF, HIT, RX, RXG, RXS, SCAN, and OCC, may
be used with the DISPLAY command to display the record for a specified
Accession Number.

ENTER DISPLAY FORMAT (FCRDREF):ed
'ED' IS NOT A VALID FORMAT FOR FILE 'CASREACT'

The following are valid formats:

ABS ----- GI and AB
ALL ----- BIB, AB, IND, RE, Single-step Reactions
APPS ----- AI, PRAI
BIB ----- AN, plus Bibliographic Data
CAN ----- List of CA abstract numbers without answer numbers
CBIB ----- AN, plus Compressed Bibliographic Data
DALL ----- ALL, delimited (end of each field identified)
IABS ----- ABS, indented with text labels
IALL ----- ALL, indented with text labels
IBIB ----- BIB, indented with text labels
IND ----- Indexing data

stn

IPC ----- International Patent Classifications
ISTD ----- STD, indented with text labels
OBIB ----- AN, plus Bibliographic Data (original)
OIBIB ----- OBIB, indented with text labels

SBIB ----- BIB, no citations
SIBIB ----- IBIB, no citations

MAX ----- Same as ALL
PATS ----- PI, SO
SCAN ----- TI and FCRD (random display, no answer number. SCAN
must be entered on the same line as DISPLAY, e.g.,
D SCAN.)
SSRX ----- Single-Step Reactions (Map, Diagram, and Summary for
all single-step reactions)
STD ----- BIB, IPC, and NCL

CRD ----- Compact Display of All Hit Reactions
CRDREF ----- Compact Reaction Display and SO, PY for Reference
FHIT ----- Reaction Map, Diagram, and Summary for first
hit reaction
FHITCBIB --- FHIT, AN plus CBIB
FCRD ----- First hit in Compact Reaction Display (CRD) format
FCRDREF ---- First hit in Compact Reaction Display (CRD) format with
CA reference information (SO, PY). (Default)
FPATH ----- PATH, plus Reaction Summary for the "long path"
FSPATH ----- SPATH, plus Reaction Summary for the "short path"
HIT ----- Reaction Map, Reaction Diagram, and Reaction
Summary for all hit reactions and fields containing
hit terms
OCC ----- All hit fields and the number of occurrences of the
hit terms in each field. Includes total number of
HIT, PATH, SPATH reactions. Labels reactions that have
incomplete verifications.
PATH ----- Reaction Map and Reaction Diagram for the "long
path". Displays all hit reactions, except those
whose steps are totally included within another hit
reaction which is displayed
RX ----- Hit Reactions (Map, Diagram, Summary for all hit reactions)
RXG ----- Hit Reaction Graphics (Map and Diagram for all hit reactions)
RXL ----- Hit Reaction Long (Map, Diagram, Summary for all hit reactions)
RXS ----- Hit Reaction Summaries (Map and Summary for all hit reactions)
SPATH ----- Reaction Map and Reaction Diagram for the "short
path". Displays all single step reactions which
contain a hit substance. Also displays those
multistep reactions that have a hit substance in both
the first and last steps of the reaction, except for
those hit reactions whose steps are totally included
within another hit reaction which is displayed

To display a particular field or fields, enter the display field
codes. For a list of the display field codes, enter HELP DFIELDS
at an arrow prompt (=>). Examples of combinations include: D TI;
D BIB RX; D TI, AU, FCRD. The information is displayed in the same order
as the specification. All of the formats, except CRD, CRDREF, FHIT, PATH,
FPATH, SPATH, FSPATH, FCRD, FCRDREF, HIT, RX, RXG, RXS, SCAN, and OCC, may

stn

be used with the DISPLAY command to display the record for a specified Accession Number.

ENTER DISPLAY FORMAT (FCRDREF):end

=> d his

(FILE 'HOME' ENTERED AT 06:48:32 ON 27 OCT 2008)


FILE 'CASREACT' ENTERED AT 06:48:45 ON 27 OCT 2008

L1 STRUCTURE UPLOADED
L2 0 S L1, CRD IBIB ABS, FHIT,
L3 0 S L1
L4 23 S L1 FULL

=> d l4, fhit ibib abs, 1-23

L4 ANSWER 1 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(19) OF 22 AP ==> AQ



RX(19) RCT AP 6298-19-7

STAGE(1)

RGT C 104-15-4 TsOH
SOL 75-05-8 MeCN
CON room temperature -> 10 deg C

STAGE(2)

RGT D 7681-11-0 KI, E 7632-00-0 NaNO2
SOL 7732-18-5 Water
CON SUBSTAGE(1) 10 minutes, 10 - 15 deg C
SUBSTAGE(2) 10 deg C -> 20 deg C
SUBSTAGE(3) 50 minutes, 20 deg C

PRO AQ 78607-36-0

ACCESSION NUMBER: 146:337774 CASREACT

TITLE: A new, one-step, effective protocol for the iodination of aromatic and heterocyclic compounds via aprotic diazotization of amines

AUTHOR(S): Krasnokutskaya, Elena A.; Semenischeva, Nadya I.; Filimonov, Victor D.; Knochel, Paul

Updated Search

stn

CORPORATE SOURCE: Department of Organic Chemistry, Tomsk Polytechnic University, Tomsk, 634050, Russia
SOURCE: Synthesis (2007), (1), 81-84
CODEN: SYNTBF; ISSN: 0039-7881
PUBLISHER: Georg Thieme Verlag
DOCUMENT TYPE: Journal
LANGUAGE: English



AB We have developed a convenient one-step preparation of aromatic and some heterocyclic iodides by the sequential diazotization-iodination of the aromatic amines with a $KI/NaNO_2/p$ -TsOH system in acetonitrile at room temperature

This method has general character and allows aryl iodides with either donor or acceptor substituents in various positions to be obtained from the corresponding amines in 50-90% yield.

REFERENCE COUNT: 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 2 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(4) OF 39 ...J + N ==> O...



O
YIELD 84%

RX(4) RCT J 6226-46-6

Updated Search

stn

STAGE(1)

RGT P 7632-00-0 NaNO₂, Q 7664-39-3 HF
SOL 7664-39-3 HF
CON SUBSTAGE(1) 5 minutes, room temperature
SUBSTAGE(2) 90 minutes, room temperature

STAGE(2)


RCT N 108-67-8
CON room temperature

PRO O 322641-70-3

ACCESSION NUMBER: 145:188836 CASREACT
TITLE: Nucleophilic substitution in
tetrafluoro-4-nitropyridine derivatives and the
corresponding fluorinated diazepines: HPLC resolution
of their isomers
AUTHOR(S): Sekhri, Lakhdar
CORPORATE SOURCE: Institut de Chimie Industriel, Universite de Ouargla,
Ouargla, 30000, Algeria
SOURCE: Asian Journal of Chemistry (2005), 17(3), 1747-1766
CODEN: AJCHEW; ISSN: 0970-7077
PUBLISHER: Asian Journal of Chemistry
DOCUMENT TYPE: Journal
LANGUAGE: English
AB Tetrafluoro-4-nitropyridine derivs. have been synthesized and separated
successfully by HPLC. The resulting fluorinated amines and
4-amino-3-chlorotrifluoropyridine have also been diazotized and the
resulting diazonium ions coupled to mesitylene giving the corresponding
azo-compds. Treatment of these azo-compds. with sodium methoxide gave the
corresponding methoxy(aryldiazo)perfluoropyridines. The thermolysis of the
synthesized azo-compds. gave the corresponding diazepines in good yields.
The structural diazepine-isomers were separated by HPLC.
REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 3 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(102) OF 668 GF ==> GG...



Updated Search

stn

RX(102) RCT GF 90902-84-4

STAGE(1)

RGT FZ 7647-01-0 HCl, GH 7632-00-0 NaNO₂
SOL 7732-18-5 Water
CON SUBSTAGE(1) -5 deg C
SUBSTAGE(2) 30 minutes, -5 deg C

STAGE(2)

RGT GI 16940-81-1 H+ [PF₆]-
SOL 7732-18-5 Water
CON SUBSTAGE(1) 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C
SUBSTAGE(3) 90 deg C
SUBSTAGE(4) 90 deg C -> room temperature

STAGE(3)

RGT CI 144-55-8 NaHCO₃
SOL 7732-18-5 Water
CON room temperature, basify

PRO GG 156772-60-0

NTE petroleum ether solvent used in 2nd stage

ACCESSION NUMBER: 143:422040 CASREACT
TITLE: Diarylalkyne compounds with MCH-receptor antagonistic activity, their preparation, pharmaceutical compositions, and use in therapy

PATENT ASSIGNEE(S): Boehringer Ingelheim International GmbH, Germany
SOURCE: U.S. Pat. Appl. Publ., 62 pp.
CODEN: USXXCO

DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
US 20050239826	A1	20051027	US 2005-104915	20050413
DE 102004017935	A1	20051103	DE 2004-10200401793520040414	
CA 2559021	A1	20051103	CA 2005-2559021	20050408
WO 2005103031	A1	20051103	WO 2005-EP3683	20050408
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1740572	A1	20070110	EP 2005-716558	20050408
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR			

stn

JP 2007532593 T 20071115
PRIORITY APPLN. INFO.:

JP 2007-507706 20050408
DE 2004-10200401/793520040414
US 2004-563677P 20040420
WO 2005-EP3683 20050408

OTHER SOURCE(S): MARPAT 143:422040
GI


* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The invention relates to alkyne compds. of general formula I, which are antagonists of melanin-concentrating hormone (MCH) receptors. In compds. I, R1 is selected from C3-6 alkenyl, C3-6 alkynyl, (hydroxy-C3-7 cycloalkyl)-C1-3 alkyl, oxa-C4-7 cycloalkyl, and dihydroxy-C3-7 alkyl, each optionally substituted; R2 is independently selected from H, (un)substituted C1-8 alkyl, (un)substituted C3-7 cycloalkyl, (un)substituted Ph, (un)substituted pyridinyl, etc., or R1 and R2, together with the N atom to which they are bound, form an (un)substituted heterocycle; X is (un)substituted C1-4 alkylene; W and Z are each independently a bond or a C1-2 alkylene; Y and A are each independently (un)substituted Ph, (un)substituted pyridinyl, (un)substituted pyrimidinyl, (un)substituted pyrazinyl, etc.; B is (un)substituted C1-6 alkyl, (un)substituted C2-6 alkenyl, (un)substituted C3-7 cycloalkyl, (un)substituted Ph, (un)substituted pyridinyl, etc.; including tautomers, enantiomers, salts, and mixts. thereof, with 6 specific compds. excluded. The invention also relates to the preparation of I, pharmaceutical compns. containing I and one or more physiol. acceptable excipients, inert carriers or diluents, as well as to the use of the compns. for the treatment of metabolic disorders and/or eating disorders, particularly obesity and diabetes. N-Alkylation of 3-methylpyridine with benzyl chloride followed by hydride reduction, asym. dihydroxylation, and debenzoylation gave optically active piperidinediol II. 2-Bromoethanol underwent substitution with 4-iodo-2-methylphenol to give the corresponding ether, which was coupled with trimethylsilylacetylene and desilylated to give alkyne III. Coupling of III with 2,5-dibromopyridine, Suzuki coupling with 4-chlorophenylboronic acid, mesylation and substitution with piperidinediol II resulted in the formation of diarylalkyne IV. The compds. of the invention are MCH-receptor antagonists, with compound IV expressing an IC50 value of 10.9 nM.

L4 ANSWER 4 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(114) OF 747 KA ==> KB...

stn



RX(114) RCT KA 90902-84-4

STAGE(1)

RGT FQ 7647-01-0 HCl, DM 7632-00-0 NaNO2
SOL 7732-18-5 Water
CON 30 minutes, -5 deg C

STAGE(2)

RGT KC 16940-81-1 H+ [PF6]-
SOL 7732-18-5 Water
CON 1 hour, 0 deg C

STAGE(3)

RGT DF 497-19-8 Na2CO3
SOL 7732-18-5 Water
CON SUBSTAGE(1) 90 deg C
SUBSTAGE(2) 90 deg C -> room temperature
SUBSTAGE(3) room temperature, basify

PRO KB 156772-60-0

NTE petroleum ether solvent used in stage 3 substage 1,
regioselective

ACCESSION NUMBER: 143:405812 CASREACT

TITLE: Preparation of substituted pyridine alkynes with MCH
antagonistic activity for the treatment of metabolic
disorders

INVENTOR(S): Stenkamp, Dirk; Mueller, Stephan Georg; Lustenberger,
Philipp; Lehmann-Lintz, Thorsten; Roth, Gerald
Juergen; Rudolf, Klaus; Schindler, Marcus; Thomas,
Leo; Lotz, Ralf

PATENT ASSIGNEE(S): Boehringer Ingelheim International GmbH, Germany

SOURCE: U.S. Pat. Appl. Publ., 67 pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20050234101	A1	20051020	US 2005-104889	20050413

stn

DE 102004017934	A1	20051103	DE 2004-10200401793420040414
CA 2559688	A1	20051103	CA 2005-2559688 20050408
WO 2005103002	A2	20051103	WO 2005-EP3685 20050408
WO 2005103002	A3	20060202	


W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

EP 1737823	A2	20070103	EP 2005-737015 20050408
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR			
JP 2007532595	T	20071115	JP 2007-507708 20050408

PRIORITY APPLN. INFO.:
 DE 2004-10200401793420040414
 US 2004-563590P 20040420
 WO 2005-EP3685 20050408

GI



II


AB Various substituted pyridinyl alkynes are prepared For instance, 2-[[4-[[5-(4-chlorophenyl)pyridin-2-yl]ethynyl]-2-methylphenyl]oxy]ethyl methanesulfonate (I) is prepared in 6 steps from 4-iodophenol, 2-bromoethanol, trimethylsilylacetylene, 2,5-dibromopyridine and 4-chlorophenylboronic acid. This intermediate is reacted with a variety of amines to produce example compds. I is converted to II by displacement with the corresponding amine. II exhibits an IC₅₀ = 6.2 nM for MCH-1. Example compds. are useful for the treatment of metabolic disorders and/or eating disorders, particularly obesity and diabetes.

L4 ANSWER 5 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

Updated Search

stn

RX(1) OF 11 ...A ==> B



RX(1) RCT A 6298-19-7

STAGE(1)

RGT C 7647-01-0 HCl
SOL 7732-18-5 Water
CON room temperature -> -8 deg C

STAGE(2)

RGT D 7632-00-0 NaNO₂
SOL 7732-18-5 Water
CON 30 minutes, -7 - -3 deg C

STAGE(3)

RGT C 7647-01-0 HCl
CAT 1317-38-0 CuO
SOL 109-69-3 BuCl, 7732-18-5 Water
CON 55 - 62 deg C

PRO B 2402-77-9

ACCESSION NUMBER: 143:155307 CASREACT
TITLE: Process for the manufacture of 2,3-dichloropyridine
INVENTOR(S): Shapiro, Rafael
PATENT ASSIGNEE(S): E.I. Dupont de Nemours and Company, USA
SOURCE: PCT Int. Appl., 23 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005070888	A2	20050804	WO 2005-US2462	20050121
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,				

Updated Search


stn

AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
AU 2005206576	A1	20050804	AU 2005-206576 20050121
CA 2553850	A1	20050804	CA 2005-2553850 20050121
EP 1706381	A2	20061004	EP 2005-712075 20050121
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK, IS, YU			
CN 1910152	A	20070207	CN 2005-80002691 20050121
BR 2005006502	A	20070227	BR 2005-6502 20050121
JP 2007523065	T	20070816	JP 2006-551437 20050121
US 20070161797	A1	20070712	US 2006-583635 20060620
IN 2006DN03640	A	20070824	IN 2006-DN3640 20060623
MX 2006PA08208	A	20060831	MX 2006-PA8208 20060719
PRIORITY APPLN. INFO.:		US 2004-539068P	20040123
		WO 2005-US2462	20050121

AB A method for preparing 2,3-dichloropyridine is disclosed in which 3-amino-2-chloropyridine is contacted with an alkali metal nitrite in the presence of aqueous hydrochloric acid to form a diazonium salt; and the diazonium salt is subsequently decomposed in the presence of copper catalyst wherein at least about 50% of the copper is the copper(II) oxidation state.

L4 ANSWER 6 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(2) OF 115 J ==> B...



RX(2) RCT J 6298-19-7

STAGE(1)
 RGT K 7647-01-0 HCl
 SOL 7732-18-5 Water
 CON room temperature -> -5 deg C

STAGE(2)
 RGT L 7632-00-0 NaNO2
 SOL 7732-18-5 Water
 CON SUBSTAGE(1) <5 deg C
 SUBSTAGE(2) 10 minutes, <5 deg C

STAGE(3)

stn

RGT M 7681-11-0 KI
SOL 7732-18-5 Water
CON SUBSTAGE(1) -5 deg C
SUBSTAGE(2) <10 deg C
SUBSTAGE(3) 0 deg C -> room temperature

STAGE(4)


RGT N 1310-73-2 NaOH
SOL 7732-18-5 Water, 141-78-6 AcOEt
CON room temperature, pH 11

PRO B 78607-36-0
NTE workup

ACCESSION NUMBER: 141:410868 CASREACT
TITLE: Synthesis of Disubstituted
Imidazo[4,5-b]pyridin-2-ones
AUTHOR(S): Kuethe, Jeffrey T.; Wong, Audrey; Davies, Ian W.
CORPORATE SOURCE: Department of Process Research, Merck & Co., Inc.,
Rahway, NJ, 07065, USA
SOURCE: Journal of Organic Chemistry (2004), 69(22), 7752-7754
CODEN: JOCEAH; ISSN: 0022-3263
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
AB Regioselective palladium-catalyzed amination of 2-chloro-3-iodopyridine
followed by a subsequent palladium-catalyzed amination leads to
2,3-diaminopyridines. Treatment with triphosgene affords highly
functionalized unsym. imidazo[4,5-b]pyridin-2-ones in just three synthetic
steps. A two-step synthesis of pseudosym. disubstituted
imidazo[4,5-b]pyridin-2-ones, 1,4-disubstituted
pyrido[2,3-b]pyrazinediones, and 1,3-disubstituted
thiadiazolo[3,4-b]pyridin-2-ones is also described.
REFERENCE COUNT: 49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 7 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(1) OF 23 A ==> B...



RX(1) RCT A 6298-19-7

STAGE(1)

Updated Search

stn

RGT C 16872-11-0 HBF₄
SOL 7732-18-5 Water, 64-17-5 EtOH
CON 15 minutes, -5 deg C

STAGE(2)

RGT D 110-46-3 Isoamyl nitrite
CON SUBSTAGE(1) 5 minutes, <0 deg C
SUBSTAGE(2) 30 minutes, <0 deg C

STAGE(3)

SOL 142-82-5 Heptane
CON SUBSTAGE(1) 2 hours, <0 deg C -> reflux
SUBSTAGE(2) reflux -> 0 deg C

STAGE(4)

RGT E 1310-73-2 NaOH
SOL 7732-18-5 Water
CON 0 deg C -> room temperature


PRO B 17282-04-1

NTE thermal in stage 3

ACCESSION NUMBER: 140:423556 CASREACT
TITLE: Synthesis of 2-alkylamino-3-fluoropyridines using
Buchwald conditions
AUTHOR(S): Munson, Peter M.; Thompson, Wayne J.
CORPORATE SOURCE: Department of Medicinal Chemistry, Merck Research
Laboratories, West Point, PA, 19486, USA
SOURCE: Synthetic Communications (2004), 34(5), 759-766
CODEN: SYNCAV; ISSN: 0039-7911
PUBLISHER: Marcel Dekker, Inc.
DOCUMENT TYPE: Journal
LANGUAGE: English
AB Synthesis of 2-alkylamino-3-fluoropyridines from 2-chloro-3-fluoropyridine
using palladium-catalyzed coupling reaction under Buchwald conditions is
described.
REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 8 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(2) OF 44 H ==> I...




RX(2) RCT H 6298-19-7

Updated Search

stn

RGT J 7647-01-0 HCl, K 7681-11-0 KI, L 7632-00-0 NaNO2
PRO I 78607-36-0
SOL 7732-18-5 Water

ACCESSION NUMBER: 140:93833 CASREACT
TITLE: An Efficient Two-Step Total Synthesis of the
Quaterpyridine Nemertelline
AUTHOR(S): Bouillon, Alexandre; Voisin, Anne Sophie; Robic,
Audrey; Lancelot, Jean-Charles; Collot, Valerie;
Rault, Sylvain
CORPORATE SOURCE: UFR des Sciences Pharmaceutiques, Centre d'Etudes et de
Recherche sur le Medicament de Normandie, Caen, 14032,
Fr.
SOURCE: Journal of Organic Chemistry (2003), 68(26),
10178-10180
CODEN: JOCEAH; ISSN: 0022-3263
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
GI




AB Regioselective and univocal Suzuki cross-coupling reactions performed on halopyridinyl boronic acids provide a flexible and versatile route to a multigram scale synthesis of 2,2'-dichloro-3,4'-bipyridine (I), which allows couplings with excess pyridin-3-yl boronic acid to give a new and efficient two-step rapid synthesis of nemertelline (II), the quaterpyridine neurotoxin isolated from a Hoplonemertine sea worm.
REFERENCE COUNT: 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 9 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(1) OF 95 ...A ==> B

stn



RX(1) RCT A 426462-05-7

STAGE(1)

RGT C 32001-55-1 4-MeOC6H4CPh2Cl
SOL 7732-18-5 Water

STAGE(2)

RGT D 7632-00-0 NaNO2

STAGE(3)


RGT E 1336-21-6 NH4OH
SOL 7732-18-5 Water

PRO B 426460-72-2

ACCESSION NUMBER: 137:353202 CASREACT
TITLE: Synthesis, Nicotinic Acetylcholine Receptor Binding,
and Antinociceptive Properties of
2-exo-2-(2',3'-Disubstituted
5'-pyridinyl)-7-azabicyclo[2.2.1]heptanes: Epibatidine
Analogues
AUTHOR(S): Carroll, F. Ivy; Lee, Jeffrey R.; Navarro, Hernan A.;
Ma, Wei; Brieady, Lawrence E.; Abraham, Philip;
Damaj, M. I.; Martin, Billy R.
CORPORATE SOURCE: Chemistry and Life Sciences, Research Triangle
Institute, Research Triangle Park, NC, 27709, USA
SOURCE: Journal of Medicinal Chemistry (2002), 45(21),
4755-4761
CODEN: JMCMAR; ISSN: 0022-2623
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
GI

Updated Search

stn




I

AB A number of 2',3'-disubstituted epibatidine analogs were synthesized and evaluated in vitro for potency at nicotinic acetylcholine receptors (nAChRs) and in vivo for antinociception activity in the tail-flick and hot-plate models of acute pain and for their ability to affect core body temperature. Compds. that possessed electron-withdrawing groups (F, Cl, Br, and I) in both the 2'- and the 3'-positions showed affinities at the nAChR similar to epibatidine. However, in vivo efficacy did not correlate with affinity. 2-Exo-(3'-Amino-2'-chloro-5'-pyridinyl)-7-azabicyclo[2.2.1]heptane (I), an epibatidine analog possessing an electron-releasing amino group in the 3'-position, produced the highest affinity. Compound I was also the most selective epibatidine analog with a K_i of 0.001 nM at $\alpha\beta$ nAChRs, which is 26 times greater than that of epibatidine, and a $\alpha\beta/\alpha\gamma$ K_i ratio of 14 000, twice that of epibatidine. In vivo testing revealed that this compound potentially inhibited nicotine-induced antinociception with AD_{50} values below 1 $\mu\text{g/kg}$. Surprisingly, this same compound was also an agonist at higher doses (ED_{50} .apprx.20 $\mu\text{g/kg}$). Thus, the addition of the 3'-amino group to epibatidine confers potent antagonist activity to the compound with little effect on agonist activity. 2,3-Disubstituted epibatidine analogs possessing a 2'-amino group combined with a 3'-bromo or 3'-iodo group showed in vitro and in vivo nAChR properties similar to nicotine.


REFERENCE COUNT: 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 10 OF 23 CASREACT COPYRIGHT 2008 ACS on STN


RX(26) OF 450 ...3 AR + 2 AM ==> AS + AT...



2 AR




AR




AM

Updated Search


stn



AM



AS
YIELD 47%



AT
YIELD 28%

RX(26) RCT AR 14036-06-7, AM 6256-96-8
PRO AS 405230-95-7, AT 405231-20-1

ACCESSION NUMBER: 136:263363 CASREACT

TITLE: Synthesis of halogen-substituted 3-deazaadenosine and 3-deazaguanosine analogues as potential antitumor/antiviral agents

AUTHOR(S): Liu, Mao-Chin; Luo, Mei-Zhen; Mozdziessz, Diane E.; Lin, Tai-Shun; Dutschman, Ginger E.; Gullen, Elizabeth A.; Cheng, Yung-Chi; Sartorelli, Alan C.

CORPORATE SOURCE: Department of Pharmacology and Developmental Therapeutics Program, Cancer Center, Yale University School of Medicine, New Haven, CT, 06520-8066, USA

SOURCE: Nucleosides, Nucleotides & Nucleic Acids (2001), 20(12), 1975-2000

PUBLISHER: CODEN: NNAFY; ISSN: 1525-7770
Marcel Dekker, Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Various 2-halogen-substituted analogs, 3-halogen-substituted analogs, and 2',3'-dihalogen-substituted analogs of 3-deazaadenosine and 3-halogen-substituted analogs of 3-deazaguanosine have been synthesized as potential anticancer and/or antiviral agents. Among these compds., 3-deaza-3-bromoguanosine showed significant cytotoxicity against L1210, P388, CCRF-CEM and B16F10 cell lines in vitro, producing IC50 values of 3, 7, 9 and 7 μ M, resp. Several 3-deazaadenosine analogs showed moderate


stn

to weak activity against hepatitis B virus.

REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 11 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(24) OF 51 AU ==> B...



RX(24) RCT AU 6298-19-7

STAGE(1)

RGT L 7647-01-0 HCl, AV 7632-00-0 NaNO2
SOL 7732-18-5 Water

STAGE(2)

RGT AW 7681-11-0 KI
SOL 7732-18-5 Water

PRO B 78607-36-0

ACCESSION NUMBER: 136:216632 CASREACT

TITLE: Coupling Reaction of Zirconacyclopentadienes with Dihalonaphthalenes and Dihalopyridines: A New Procedure for the Preparation of Substituted Anthracenes, Quinolines, and Isoquinolines

AUTHOR(S): Takahashi, Tamotsu; Li, Yanzhong; Stepnicka, Petr; Kitamura, Masanori; Liu, Yanzun; Nakajima, Kiyohiko; Kotora, Martin

CORPORATE SOURCE: Catalysis Research Center and Graduate School of Pharmaceutical Sciences, Hokkaido University, Japan
SOURCE: Journal of the American Chemical Society (2002), 124(4), 576-582

CODEN: JACSAT; ISSN: 0002-7863

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Reactions of tetraiodobenzene with zirconacyclopentadienes, which were conveniently prepared from two alkynes (or diynes) and zirconocene complexes, afforded 1,2,3,4-tetrasubstituted diiodonaphthalene derivs. in good isolated yields. These 1,2,3,4-tetrasubstituted diiodonaphthalene derivs. could be converted to 1,2,3,4,5,6,7,8-octasubstituted anthracene derivs. by reaction with a second zirconacyclopentadiene. When the two zirconacyclopentadienes were different, unsym. anthracenes such as

Updated Search


stn

1,2,3,4-tetraethyl-5,6,7,8-tetraphenylanthracene (68% isolated yield) were obtained. On the other hand, treatment of a 2,3-dihalopyridine such as 2-bromo-3-iodopyridine with zirconacyclopentadienes gave 5,6,7,8-tetrasubstituted quinoline derivs. in good to high yields. 3,4-Dihalopyridines such as 4-chloro-3-iodopyridine reacted with zirconacyclopentadienes to afford 5,6,7,8-tetrasubstituted isoquinoline derivs. in good to high yields.

REFERENCE COUNT: 55 THERE ARE 55 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 12 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(5) OF 30 ...F ==> L...



RX(5) RCT F 39745-40-9
RGT J 7632-00-0 NaNO₂, K 7647-01-0 HCl
PRO L 54957-86-7
NTE 5-10.deg., CHLORIDES

ACCESSION NUMBER: 126:47080 CASREACT
TITLE: Synthesis of dihalopicoline N-oxides and their 4-nitro derivatives

AUTHOR(S): Ciurla, H.; Puszko, A.

CORPORATE SOURCE: Russia

SOURCE: Khimiya Geterotsiklicheskikh Soedinenii (1996), (10), 1366-1371

CODEN: KGSSAQ; ISSN: 0132-6244

PUBLISHER: Latviiskii Institut Organicheskogo Sintez

DOCUMENT TYPE: Journal


LANGUAGE: English

AB Three aminohalo-substituted α - and β -picolines, six dihalo-substituted α - and β -picolines, six dihalo-substituted α - and β -picoline N-oxides, and six dihalo-4-nitropicoline N-oxides were synthesized in excellent yields. Some properties of the products were reported.

L4 ANSWER 13 OF 23 CASREACT COPYRIGHT 2008 ACS on STN


RX(1) OF 17 A ==> B

STN



RX(1) RCT A 6298-19-7
RGT C 7632-00-0 NaNO₂, D 7787-70-4 CuBr, E 10035-10-6 HBr
PRO B 52200-48-3
SOL 108-88-3 PhMe

ACCESSION NUMBER: 122:186872 CASREACT
TITLE: Use of Hydrogen Bonds to Control Molecular Aggregation. Behavior of Dipyridones and Pyridone-Pyrimidones Designed To Form Cyclic Triplexes
AUTHOR(S): Boucher, Eric; Simard, Michel; Wuest, James D.
CORPORATE SOURCE: Departement de Chimie, Universite de Montreal, Montreal, QC, H3C 3J7, Can.
SOURCE: Journal of Organic Chemistry (1995), 60(5), 1408-12
CODEN: JOCEAH; ISSN: 0022-3263
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
GI




AB The tendency of 2-pyridones and related heterocycles to form cyclic hydrogen-bonded dimers allows them to be used as sticky sites that induce mols. in which they are incorporated to associate in particular ways. I, which is constructed from pyridone and pyrimidone subunits linked to a rigid linear acetylenic spacer, incorporates an array of hydrogen-bonding sites designed to favor the formation of a cyclic triplex. I was prepared and the structure of its DMSO solvate was determined by X-ray crystallog. Aggregation does not produce a cyclic triplex but rather gives chains in which adjacent mols. of I are linked by single hydrogen bonds.

L4 ANSWER 14 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(14) OF 66 COMPOSED OF RX(4), RX(5)
RX(14) J ==> R

Updated Search

stn



RX(4) RCT J 154012-16-5
RGT O 16940-81-1 H+ [PF6]-, P 7632-00-0 NaNO2
PRO N 154012-09-6
SOL 7732-18-5 Water

RX(5) RCT N 154012-09-6
PRO R 154012-17-6
NTE thermal; key step



ACCESSION NUMBER: 120:244961 CASREACT

TITLE: The synthesis of a series of
7-amino-1-cyclopropyl-8-fluoro-1,4-dihydro-4-oxo-1,6-
naphthyridine-3-carboxylic acids as potential
antibacterial agents

AUTHOR(S): Sanchez, Joseph P.; Gogliotti, Rocco D.
CORPORATE SOURCE: Parke-Davis Pharm. Res. Div., Warner-Lambert Co., Ann
Arbor, MI, 48105, USA

SOURCE: Journal of Heterocyclic Chemistry (1993), 30(4), 855-9
CODEN: JHTCAD; ISSN: 0022-152X

DOCUMENT TYPE: Journal
LANGUAGE: English
GI




AB A series of title compds. I [R = 3-aminopyrrolidin-1-yl, 3-(ethylaminomethyl)pyrrolidin-1-yl, 4-aminopiperidin-1-yl, piperazin-1-yl] was prepared and evaluated for antibacterial activity (no data). I were prepared by the displacement of the chloro substituent from I (R = Cl) with the requisite nitrogen nucleophile. The naphthyridine acid was synthesized in ten steps from pyridinecarboxylate II (R1 = OH, R2 = NO2). The key step in the sequence was a Schiemann reaction of II (R1 =

stn

Cl, R2 = N2+ PF6-) to give II (R1 = Cl, R2 = F).


L4 ANSWER 15 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(2) OF 4 ...B ==> H



RX(2) RCT B 152840-65-8
RGT I 7787-70-4 CuBr, J 10035-10-6 HBr
PRO H 152840-66-9
NIE NANO

ACCESSION NUMBER: 120:107714 CASREACT
TITLE: A synthetic approach to carbon-14 labeled
antibacterial naphthyridine- and quinolonecarboxylic
acids
AUTHOR(S): Ekhatu, I. Victor; Huang, Che C.
CORPORATE SOURCE: Parke-Davis Pharm. Res., Warner-Lambert Co., Ann
Arbor, MI, 48105, USA
SOURCE: Journal of Labelled Compounds and Radiopharmaceuticals
(1993), 33(9), 869-80
CODEN: JLCRD4; ISSN: 0362-4803
DOCUMENT TYPE: Journal
LANGUAGE: English
GI



AB Labeled versions of (S)-clinafloxacin (I; R = H, X = CCl) and two
naphthyridinecarboxylic acid antibacterial compds. (I; R = H, H-Ala, X =
N) were prepared. Preps. started from hitherto unknown bromo compds. II (R1
= Br), from which the corresponding 14C-labeled aromatic carboxylic acids II


Updated Search

stn

(R = 14CO2H) were generated by metal-halogen exchange followed by carboxylation reaction. Details of these preps. are given.

L4 ANSWER 16 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(24) OF 121 B + AU ==> AV




AV
YIELD 35%

RX(24) RCT B 1572-52-7, AU 55304-76-2
RGT D 7647-01-0 HCl, E 110-46-3 Isoamyl nitrite
PRO AV 112177-06-7
CAT 7758-89-6 CuCl
SOL 756-79-6 MeP(O)(OMe)2

ACCESSION NUMBER: 108:55848 CASREACT
TITLE: The synthesis of halogenated pyridines substituted at the carbon atom C-3
AUTHOR(S): Sutter, Peter; Weis, Claus D.
CORPORATE SOURCE: Dyest. Chem. Dep., Ciba-Geigy, Ltd., Basel, Switz.
SOURCE: Journal of Heterocyclic Chemistry (1987), 24(4), 1093-102
CODEN: JHTCAD; ISSN: 0022-152X
DOCUMENT TYPE: Journal
LANGUAGE: English
GI

Updated Search


stn



AB Seventeen 3-substituted pyridines I (R = Ph, 4-MeC6H4, 4-NO2C6H4, 2,5-Cl2C6H3, 3-pyridinyl, etc.) were prepared in 3 steps from the corresponding amines RNH2 (II). Arylation of H2C:C(CN)CH2CH2CN with II in the presence of CuCl, HCl, and isoamyl nitrite in di-Me methylphosphonate (preferred solvent) gave dicyanobutanes RCH2CCl(CN)CH2CH2CN which were cyclized with H2SO4-HOAc to give piperidinediones III. Aromatization with POCl3 in the presence of HMPA gave I.

L4 ANSWER 17 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(21) OF 90 ...AO ==> AR...




RX(21) RCT AO 109613-97-0
RGT AS 11113-50-1 Boric acid, AT 7664-39-3 HF, AU 110-46-3 Isoamyl nitrite
PRO AR 109613-98-1
SOL 7732-18-5 Water, 64-17-5 EtOH
NTE thermal diazonium salt decompn. in ligroin

ACCESSION NUMBER: 107:198450 CASREACT
TITLE: Syntheses of hydroxylated 2,2'-bipyridines. I. Orellanine, the poison of a toadstool
AUTHOR(S): Dehmlow, Eckehard V.; Schulz, Hans Joachim
CORPORATE SOURCE: Fak. Chem., Univ. Bielefeld, Bielefeld, D-4800, Fed. Rep. Ger.
SOURCE: Liebigs Annalen der Chemie (1987), (10), 857-61
CODEN: LACHDL; ISSN: 0170-2041
DOCUMENT TYPE: Journal
LANGUAGE: German
GI

Updated Search


stn



AB Orellanine (I) was prepared from 2-chloro-3-fluoropyridine or 3-amino-4-methoxypyridine via biaryl coupling of 2-chloro-3,4-dimethoxypyridine or 2-bromo-3-fluoro-4-methoxypyridine, resp. Reaction of I with CH₂N₂ gave bipyridines II and III. Results of UV irradiation of I are also given.

L4 ANSWER 18 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(35) OF 76 COMPOSED OF RX(9), RX(10)
 RX(35) R + S ==> D



RX(9) RCT R 58596-89-7, S 124-40-3
 RGT U 7647-01-0 HCl, V 7632-00-0 NaNO₂
 PRO T 104866-47-9

RX(10) RCT T 104866-47-9
 RGT W 7664-39-3 HF

Updated Search

stn

PRO D 104866-49-1

ACCESSION NUMBER: 105:191059 CASREACT
 TITLE: 1-Cyclopropyl-1,4-dihydro-4-oxo-1,8-naphthyridine-3-carboxylic acids
 INVENTOR(S): Petersen, Uwe; Grohe, Klaus; Zeiler, Hans Joachim; Metzger, Karl Georg
 PATENT ASSIGNEE(S): Bayer A.-G. , Fed. Rep. Ger.
 SOURCE: Ger. Offen., 64 pp.
 CODEN: GWXXBX
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:



PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 3508816	A1	19860710	DE 1985-3508816	19850313
NO 8505134	A	19860711	NO 1985-5134	19851218
NO 163331	B	19900129		
NO 163331	C	19900509		
EP 187376	A2	19860716	EP 1985-116551	19851224
EP 187376	A3	19880504		
EP 187376	B1	19920513		
R: AT, BE, CH, DE, FR, GB, IT, LI, NL, SE				
AT 76076	T	19920515	AT 1985-116551	19851224
US 4840954	A	19890620	US 1985-815440	19851231
IL 77538	A	19920525	IL 1986-77538	19860107
FI 8600073	A	19860711	FI 1986-73	19860108
FI 86721	B	19920630		
FI 86721	C	19921012		
DD 241258	A5	19861203	DD 1986-286039	19860108
DD 257427	A5	19880615	DD 1986-296482	19860108
DD 257428	A5	19880615	DD 1986-296483	19860108
CA 1339373	C	19970826	CA 1986-499241	19860108
DK 8600091	A	19860711	DK 1986-91	19860109
DK 168439	B1	19940328		
JP 61161284	A	19860721	JP 1986-1485	19860109
JP 06053741	B	19940720		
ZA 8600163	A	19860924	ZA 1986-163	19860109
HU 40126	A2	19861128	HU 1986-87	19860109
HU 193623	B	19871130		
AU 8652164	A	19870122	AU 1986-52164	19860109
AU 574550	B2	19880707		
ES 550767	A5	19880715	ES 1986-550767	19860109
PL 148191	B1	19890930	PL 1986-264565	19860109
PL 148759	B1	19891130	PL 1986-257419	19860109
HU 202840	B	19910429	HU 1987-1847	19860109
CN 86100126	A	19860709	CN 1986-100126	19860110
CN 1003239	B	19890208		
NO 8600199	A	19860711	NO 1986-199	19860121
AU 8773118	A	19870910	AU 1987-73118	19870515
AU 576449	B2	19880825		
AU 8818359	A	19880915	AU 1988-18359	19880624
FI 8902675	A	19890601	FI 1989-2675	19890601
CA 1320206	C2	19930713	CA 1990-615694	19900405
PRIORITY APPLN. INFO.:			DE 1985-3500562	19850110

Updated Search

stn

DE 1985-3508816 19850313
EP 1985-116551 19851224
CA 1986-499241 19860108
FI 1986-73 19860108


OTHER SOURCE(S): MARPAT 105:191059
GI



AB The title compds. [I; R = halo, NO₂; R₁ = (un)substituted 1-piperazinyl, 1-pyrrolidinyl] were prepared as bactericides and feed additives. Thus, 2,6-dichloro-5-methyl-3-pyridinamine (II, R₂ = NH₂, R₃ = Me) was diazotized and coupled with Me₂NH to give II (R₂ = Me₂NN, R₃ = Me) which was fluorinated with HF to give II (R₂ = F, R₃ = Me). The latter was converted in 6 steps to II (R₂ = F, R₃ = EtO₂CC(:CHOEt)CO) which was condensed with cyclopropylamine, followed by cyclization and hydrolysis of the ester group, to give I (R = F, R₁ = Cl). The latter was heated with piperazine in Me₂SO to give I (R = F, R₁ = 1-piperazinyl) (III). III had a min. inhibitory concentration of ≤0.015 mcg/mL against Escherichia coli Neum. Tablets were prepared each containing III 583.0, microcryst. cellulose 55.0, cornstarch 72.0, polyvinylpyrrolidone 30.0, dispersed silica 5.0, and Mg stearate 5.0 mg.

L4 ANSWER 19 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(19) OF 49 ...J ==> AO...



RX(19) RCT J 104829-98-3

STAGE(1)

RGT AQ 7632-00-0 NaNO₂, AR 7647-01-0 HCl

Updated Search

stn

SOL 7732-18-5 Water

STAGE(2)

RGT AS 7681-11-0 KI

SOL 7732-18-5 Water

PRO AO 104830-09-3

ACCESSION NUMBER: 105:172323 CASREACT

TITLE: Condensed heteroaromatic ring systems. IV. Synthesis of naphthyridine derivatives by cyclization of aminopyridineacrylic esters

AUTHOR(S): Sakamoto, Takao; Kondo, Yoshinori; Yamanaka, Hiroshi

CORPORATE SOURCE: Pharm. Inst., Tohoku Univ., Sendai, 980, Japan


SOURCE: Chemical & Pharmaceutical Bulletin (1985), 33(11), 4764-8

CODEN: CPBTAL; ISSN: 0009-2363


DOCUMENT TYPE: Journal

LANGUAGE: English

GI



I



II

AB The reaction of aminohalopyridines with Et acrylate in the presence of palladium(II) acetate and triarylphosphine gave Et aminopyridineacrylates, e.g., I. The cyclization of the resulting acrylates under basic conditions gave naphthyridinones having a carbostyryl-type moiety, e.g., II.

L4 ANSWER 20 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(4) OF 94 ...J ==> L...



J

L

RX(4) RCT J 6298-19-7

Updated Search


stn

STAGE(1)
RGT M 7782-77-6 HNO2

STAGE(2)
RGT N 16872-11-0 HBF4

PRO L 17282-04-1


ACCESSION NUMBER: 104:168236 CASREACT
TITLE: Synthesis of orellanine, the lethal poison of a
toadstool
AUTHOR(S): Dehmloew, Eckehard V.; Schulz, Hans Joachim
CORPORATE SOURCE: Fak. Chem., Univ. Bielefeld, Bielefeld, D-4800/1, Fed.
Rep. Ger.
SOURCE: Tetrahedron Letters (1985), 26(40), 4903-6
CODEN: TELEAY; ISSN: 0040-4039
DOCUMENT TYPE: Journal
LANGUAGE: English
GI



AB Orellanine, (I) was prepared in 10 steps from 3-aminopyridine, thus proving the identity of the natural product.

L4 ANSWER 21 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(1) OF 12 A ==> B...



RX(1) RCT A 39856-58-1
PRO B 40273-45-8
CAT 16872-11-0 HBF4
ACCESSION NUMBER: 99:195034 CASREACT
TITLE: Review on the metalation of π -deficient heteroaromatic compounds. Regioselective

Updated Search

stn


ortho-lithiation of 3-fluoropyridine: directing effects and application to synthesis of 2,3- or 3,4-disubstituted pyridines

AUTHOR(S): Marsais, Francis; Queguiner, Guy
CORPORATE SOURCE: Lab. Chim. Org. Heterocyclique, Inst. Natl. Super. Chim. Ind. Rouen, Mont Saint Aignan, 76130, Fr.
SOURCE: Tetrahedron (1983), 39(12), 2009-21
CODEN: TETRAB; ISSN: 0040-4020
DOCUMENT TYPE: Journal; General Review
LANGUAGE: English

AB Lithiation of 3-fluoropyridine is chemoselective at low temps. using butyllithium-polyamine chelates or lithium diisopropylamide. Protophilic attack by these strong bases can be directed either at the 2- or 4-position depending on the lithiation conditions. Various reaction parameters are studied: solvent, temperature, reaction time, lithium-chelating agent metalating agent. The high regioselectivity of 3-fluoropyridine lithiation is theor. discussed, in particular in terms of kinetic or thermodyn. control of the metalation. Chelation between butyllithium and 3-fluoropyridine is proposed, which completely modifies the heterocycle reactivity toward the lithiating agent. This is confirmed by theor. quantum calcns. performed on different models of 3-fluoropyridine using the CNDO/2. These results permit selection of 3-fluoropyridine metalation conditions which lead to 3-fluoro-2-lithiopyridine on the one hand and to 3-fluoro-4-lithiopyridine on the other hand. Each of the lithiated isomers is then reacted with a great variety of electrophiles to give the corresponding 2,3- or 3,4-disubstituted pyridines. Metalation of π -deficient heterocycles was also reviewed.

L4 ANSWER 22 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(3) OF 30 G ==> H...



RX(3) RCT G 6298-19-7
RGT I 7782-77-6 HNO₂, J 7787-70-4 CuBr
PRO H 52200-48-3

ACCESSION NUMBER: 88:152364 CASREACT
TITLE: Synthesis and pharmacological properties of certain alkylcarbamoylpyridinesulfonamides
AUTHOR(S): Delarge, J.
CORPORATE SOURCE: Inst. Pharm., Univ. Liege, Liege, Belg.
SOURCE: Acta Poloniae Pharmaceutica (1977), 34(3), 245-9

Updated Search

stn

CODEN: APPHAX; ISSN: 0001-6837


DOCUMENT TYPE:

Journal


LANGUAGE:

French

GI



I





II

AB Sixteen pyridine analogs I (R = 3-, 4-, 5-, 6-Me, 2-, 4-, 6-Cl, 3-Br, 4-Et2N, 4-Me2CHNH, 4-(3-ClC6H4)NH, 4-(3-CF3C6H4)NH; R1 = Et, Pr, Me2CH, Bu; SO2NHCONHR1 (in 2, 3, and 4 positions) of hypoglycemic sulfonamides were prepared from the appropriate II and R1NCO. II (R = 3-Br; SO2NH2 in 2 position) was prepared by converting 2-chloro-3-aminopyridine into 2-chloro-3-bromopyridine in a Sandmeyer reaction, then followed by reaction with KSH to give 3-bromopyridine-2-thiol, which was oxidized with Cl followed by amidation. I revealed no hypoglycemic activity; some of them were mild antiinflammatory agents. The 4-aryl-3-sulfonamide derivs. of the type I (R1 = Pr and Bu) were strong diuretics in expts. with animals as well as in clin. tests.

L4 ANSWER 23 OF 23 CASREACT COPYRIGHT 2008 ACS on STN

RX(15) OF 22 AD ==> AE



AD 

AE
YIELD 70%


RX(15) RCT AD 55304-76-2
RGT E 10035-10-6 HBr
PRO AE 55304-89-7

ACCESSION NUMBER: 84:121615 CASREACT
TITLE: Halogenated pyridines. V. Fluorinated and brominated pyridine compounds
AUTHOR(S): Mutterer, Francis; Weis, Claus D.
CORPORATE SOURCE: Div. Kunstst.-Addit. Farbst.-Chem., Ciba-Geigy A.-G., Basel, Switz.
SOURCE: Helvetica Chimica Acta (1976), 59(1), 229-35

Updated Search

DOCUMENT TYPE:
LANGUAGE:
GI

CODEN: HCACAV; ISSN: 0018-019X
Journal
German



AB Fluoropyridines I (R = F, R₁ = Cl, Me, CF₃, NO₂, R₂ = H, R₁ = Cl, Me, R₂ = Cl) were prepared by treating I (R = Cl, Br) with KF. I (R = Cl, R₁ = CF₃, R₂ = H) was obtained by treating I (R = Cl, R₁ = CCl₃, R₂ = H) with HF or SbF₃. The bromopyridines II (R₃ = Br; R₄ = H, Cl, CH₂R₃, NO₂, CHO, CO₂H, CF₃, NH₂; R₅ = H, R₃; R₆ = H, Cl, NO₂) were obtained by brominating II (R₃ = Cl) with HBr-HOAc.